**Composition**: Clopidogrel+Aspirin; (75mg+75mg) Tablet.

Indications: This tablet is indicated for the reduction of atherosclerotic events (myocardial infarction, stroke and vascular death) in patients with atherosclerosis documented by recent stroke, recent myocardial infarction or established peripheral arterial disease.

**Dosage and administration :** The recommended dose of this tablet is one tablet once daily or as directed by the registered physician.

**Contraindication :** The use of this tablet is contraindicated in the following conditions:

# Hypersensitivity to the drug substance of any component of the product.

# Hypersensitivity to Aspirin and / or NSAIDs.

# Active pathological bleeding such as peptic ulcer or intracranial hemorrhage of bleeding disorders like hemophilia. Recent history of gastrointestinal bleeding.

Side effects: Adverse effects of this tablet is minor & in rare case gastrointestinal hemorrhage, neutropenia, myelotoxicity, abdominal pain, fatigue, dyspepsia, gastritis, constipation, skin rash, headache, chest pain, diarrhoea, rhinitis, nausea dizziness, edema may occur.

**Precautions**: As with other antiplatelet agents this tablet should be used with caution in patients who may be at risk of increased bleeding from trauma, surgery

## **Clopid-As**

**Film Coated Tablet** 

or other pathological conditions.

## Use in pregnancy and lactation:

There are no adequate and well-controlled studies of Clopidogrel with Aspirin in pregnant women. However, Clopidogrel with Aspirin should be used during pregnancy only if clearly needed. This tablet should not be prescribed to nursing mother unless absolutely necessary.

Drug interactions: Clopid-As may enhance the effect of anticoagulants. No significant interaction were observed in the Co-administration with atenolol, Nifedipine or both of the drugs. Caution shold be used in Co-administration of phenytoin, tamoxifen, tolbutamide, warfarin, and fluvastatin with this tablet for possible impaired metabolism.

Packing: Clopid-As: 3x10's tablets in blister pack.