

Composition : Each Film Coated Tablet contains 1250mg Calcium Carbonate BP (Eq. to. Elemental Calcium 500mg) & Colecalciferol (Vitamin D₃) BP 200 IU.

Indications : Calcium and Vitamin D is prescribed for treatment of osteoporosis, osteomalacia, rickets, tetany, and parathyroid disease. Calcium supplements are often used to ensure adequate intake in conditions such as pregnancy, lactation, osteogenesis and tooth formation (adjunct with definite treatment) and with anti-seizure medications.

Dosage & administration : Two tablets daily or one tablet twice daily. It is best taken with or just after a meal to improve absorption. Each dose should be taken with a full glass of water. Or, as directed by the registered physician.

Contraindications : It is contraindicated in case of hypercalcaemia, hyperthyroidism, renal calculi & nephrolithiasis, Zollinger-Elison syndrome and concomitant digoxin therapy.

Precautions : If there is any pre-existing heart disease or kidney disease, precautions should be taken.

Side effects : This combination is well tolerated. Constipation may occur. If there is experience like nausea, vomiting, stomach cramps, dry mouth, increased thirst, increased urination while taking, physician should be notified.

Caldil-Plus

Tablet



DRUG
INTERNATIONAL
LTD.

Use in Pregnancy and Lactation :

Caldil-Plus should be used as directed by registered physician during pregnancy or while breast feeding.

Use in child : Caution should be taken when it is administered to a child.

Drug interactions : It has been possible interaction with digoxin, antacids containing calcium, aluminium or magnesium, other calcium supplements, calcitriol or other vitamin D supplements; tetracycline, doxycycline, aminocycline or oxytetracycline etc. So before taking any of these drugs with Caldil-Plus, suggestions of the physicians are needed.

Overdose : Assess patient for nausea, vomiting, anorexia, thirst, severe constipation, paralytic ileus and bradycardia. Contact physician or other health care professional immediately if these signs of hypercalcemia occur. □

Storage : Store below 30°C in a dry place.

Packing : Each box contains 50's tablets in blister pack.