

**Composition** : Each ml contains Polyethylene Glycol 400 BP 4mg and Propylene Glycol BP 3mg.

**Pharmacology** : The combination of Polyethylene Glycol 400 & Propylene Glycol is clinically proven to reduce both signs and symptoms of dry eye. The mechanism of action is thought to be due to its unique gelling and lubricating system formulated to adjust to each user's individual tear pH. When the ingredients of I-Moist combine with natural tears, a soft gel forms a network of protection over the eye surface. Since it promotes a healthy environment in eye surface, damaged surface cells of eye can repair more easily.

**Indications** : It is indicated for the temporary relief of burning and irritation due to dryness of the eye.

**Dosage & Administration** : Instill 1 or 2 drops in the affected eye(s) 4 times daily or as needed. Or, as directed by the registered physician.

**How to use single dose :**

- \* Wash hands before start. Open plastic pouch & remove the plastic container.
- \* Twist off the plastic container's cap.
- \* Hold the plastic container pointing down, between thumb and fingers.
- \* Tilt head back.
- \* Pull down lower eyelid with a finger, until there is a 'pocket' between the eyelid and eye. The drop will go in here.
- \* Bring the tip of the plastic container close to the eye. Do this in front of a mirror if it helps.
- \* Do not touch eye or eyelid, surrounding

**I-Moist  
Eye Drops**



**DRUG  
INTERNATIONAL  
LTD.**

areas or other surfaces with the plastic container.

\* Gently press on the base of the plastic container to release one drop at a time.

**Contraindications** : It is contraindicated in patients with known hypersensitivity to any ingredient of this formulation.

**Precautions** : Never touch tip of container with any surface to avoid contamination.

**Side effects** : Generally well tolerated. This sterile eye drops should not be used if allergic condition occurs to any ingredients of the product.

**Use in Pregnancy and Lactation** : There are no adequate and well-controlled studies in pregnant women. Caution should be exercised when administered to a nursing woman.

**Use in Child** : There is no data available.

**Drug-Interactions** : There is no data available.

**Overdose** : There is no data available.

**Storage** : Store below 30°C away from sunlight. Keep out of reach of children. Once the pouch has been opened, any unused plastic container should be stored in the open pouch. Dispose of the plastic container after use. Do not keep an open plastic container to use again.

**Packing** : Each box contains 25's (0.4ml) single use plastic containers enclosed in a plastic pouch overwrap.